

Discovery of a soft X-ray QPO from the accreting millisecond pulsar IGR J00291+5934

Ferrigno, C. Bozzo, E. Sanna, A. Pintore,
F. Papitto, A. Riggio, A. Burderi, L. Di
Salvo, T. Iaria, R. D'Ai, A.

Kulkarni & Romanova (2013)







Figure 1. *Left panel:* A 3D view of the funnel flow from the disc to a magnetized star, where the dipole moment μ is tilted by $\Theta = 20^\circ$ about the rotational axis. One of the density levels is shown in green; sample field lines are shown in red. *Right panel:* the energy flux distribution on the



Rotationally powered evolution



Accretion powered evolution

Recycled pulsars


- Accretion of material conveys angular momentum and spins-up the pulsar.

Name	P_{Spin} [ms]	P_{Orb} [min]	$M_{\text{C,Min}}$ [M_{sol}]	Discovered
SAX J1808.4-3658	2.5	120	0.043	Apr. 1998
XTE J1751-306	2.3	42	0.014	Apr. 2002
XTE J0929-314	5.4	44	0.083	Apr. 2002
XTE J1807-294	5.2	40	0.0066	Feb. 2003
XTE J1814-338	3.2	258	0.17	Jun. 2003
IGR J00291+5934	1.67	150	0.039	Dec. 2004
HETE J1900.1-2455	2.6	84	0.016	Jun. 2005
Swift J1756.9-2508	5.5	54	0.007	Jun. 2007
NGC 6440 X-2	4.86	57	0.0067	Aug. 2009
IGR J17511-3057	4.1	208	0.13	Sep. 2009
Swift J1749.4-2807	1.9	530	0.6	Apr. 2010
IGR J17498-2921	2.5	230.4	0.17	Aug 2011
IGR J18245-2452	3.9	661.5	0.17	March 2013
MAXI J0911-655	2.9	44.3	0.024	February 2016

+ 2 Intermittent pulsars: Aql X-1 and SAX J1748.9-2021

- Fastest AMSP: 600 Hz (fastest pulsar 716 Hz, Hessels et al. 2006)
- Measured spin-up in accretion phase ($8.4 \cdot 10^{-13}$ Hz/s, Falanga et al, 2005)
- Zero orbital period derivative, limit compatible with Gravitational Wave emission (Patruno et al., 2016)
- Compact binary, similar to SAX J1808.4 companion mass $0.04-0.16 M_{\text{sol}}$
- A Type-C QPO was found in 2004 outburst (Linares et al. 2007). Frequency at 44 mHz and a hard spectrum.
- In 2015 the source underwent its brightest outburst (this study) and a thermonuclear burst was detected for the first time with Swift (De Falco et al., 2016).
- Estimate the distance (4 kpc, De Falco et al., 2016).


IGR J00291+5934




Ferrigno et al.
(MNRAS 2017)

Thermonuclear
burst




- 2015: 70 ks XMM Epic timing mode 86 ks RGS
- 40 ks Contemporary Nustar
- Decay of the outburst




Courtesy A.
Sanna



- 0.5 -11 keV XMM EPIC-pn
- Four broad Lorentzians and one QPO: $f=8$ mHz $Q=9$ $rms=15\%$




- Divided in 12 energy bins with equal number of photons.
- Frequency at 8.5 mHz
- **A very soft QPO !**
- Rather high Q-factor




- At low luminosity, sometimes hard, sometimes soft.


- As for the average spectrum (Sanna et al. 2017), we use $(Tbabs * (nthComp (kT_s = 1, kT_e = 28 \text{ keV} + BlackBody (0.5 \text{ keV})))$
- We exploit hardness to guess the origin of variability.
- Softening due to additional black-body and lower seed-photon temperature in Comptonization. Almost equal power-law slope.

Hard Spectra (HR>4)



Soft Spectra (HR<2)







- Lags at QPO frequency
- Reconstruct energy-dependent QPO front from rms and lag (1.5% systematic).
- Black Body (both T and R) drives variability

$$w(E, \phi) = \mu(E) \left(1 + \sqrt{2} \sigma_1 \cos[\phi - \Phi_1(E)] \right)$$

rms
lag



mean spectrum





- Nearly sinusoidal. Only one pole is probably seen.

- Take a time series of *fractional* amplitudes of the 600 Hz spin, measured every ten seconds.
- Make a Power density spectrum: we find a QPO with the same energy dependency.



- Strong and Soft QPO at 8 mHz
- Driven mainly by a soft spectral component, identified as a black body with a few km radius and ~ 0.5 keV temperature
- Fractional pulsed amplitude is modulated by a QPO


- Is it a phenomenon inherent to

- the accretion flow, or
- thermonuclear burning, or
- ~~is it due to absorption?~~


Parameter	HR > 4	HR < 2
N_H (10^{22} cm $^{-2}$)	0.35 ± 0.02	$0.31^{+0.03}_{-0.05}$

- Absorption is irrelevant, as showed by spectra resolved in hardness or QPO phase


- NGC 6440 X-2 is a transient accreting millisecond pulsar with a 4.86 ms spin period and 57 min orbit




- Strong semi-period modulation at a frequency of ~ 1 Hz.
- PSD fitted with 3 harmonic Gaussians (1 Hz) + Lorentzian (10 Hz QPO)
- Similar to SAX J1808.4-3658




Patruno & D'Angelo (2013)



- Matter accumulates and then is accreted in cyclic fashion
- Contemporary ejections of material
- Much lower time scale (10^3 lower)




Altamirano et al. (2011)




- A very similar signature in the power spectrum of the black-hole binary H1743–322 was found in the 2010-2011 outburst using Chandra.
- They argue it is similar to the 1 Hz QPO seen in neutron star and interpreted as the movement of the disc in and out of the coronation radius.
- QPOs are harder (seen in RXTE above ~3 keV) and variable in frequency.

Watts et al.
(2012)




- Material accumulated on the NS surface undergoes catastrophic nuclear burning and the object becomes very bright.

Brightening








- IGR J17480-2446 (Terzan 5) transitioned from Atoll to Z-states. Becoming extremely bright, the Thermonuclear bursts became quasi periodic flares with recurrence of ~ 300 s.
- Our source is dimmer, spectrum is NOT BB dominated, LC is more “dipping” than flaring

- It is natural at a high accretion (\sim Eddington) and very narrow range of mass accretion rate (1%). In 4U 1636-53, they appear over a range of 50% in luminosity.
- To lower luminosity, it is argued that burning take place in a narrow equatorial zone. Heger (2007). Ashes of H burning.



- 4U 1636-53 shows QPO at 8 mHz with a very soft spectrum, interrupted by thermonuclear bursts. **Discovered in 2001 by Revnivtsev**
- Spectrum has a strong BB with $kT \sim 2$ keV
- Only in a (broad) range of luminosity



Altamirano et al. (2008)

Lyu et al. (2015)

- We found a very pronounced variability in the XMM-Newton observation of IGR J00291, reflected by the presence of a soft QPO (<3 keV, *below RXTE band*).
- This variability resembles the QPOs in 4U 1608-52, 4U1636-536, and Aql X-1, interpreted as *marginally stable nuclear burning*. However, **only one thermonuclear burst in IGR J00291** history and well apart both in flux and time from the QPO.
- **Variability is reflected in the pulsed fraction:** shape of accretion driven emission region? Change of the surface thermonuclear emission pattern leading to a modulation in pulsed fraction?
- We leave open the possibility that it is an accretion-flow mediated modulation of the accretion rate as in H1743–322 (or GRS 1915+105). However, why would it be so soft?
- Multi-messenger data will solve these degeneracies: see eXTP !