

## SPICAM UV Observations of Thermospheric Airglow and H Corona Variability

M. Chaffin (1), D. Everding (1), N. Schneider (1), F. Leblanc (2), J. Y. Chaufray (2), F. Montmessin (2), J.-L. Bertaux (2) and the SPICAM team

<sup>1</sup>Laboratory for Atmospheric and Space Physics, Boulder, Colorado, USA,


<sup>2</sup>LATMOS/IPSL, Guyancourt, France

**Introduction:** In 2007 (Mars Year 28), Mars experienced a global dust storm. Previous studies have reported large changes in the H Lyman alpha coronal airglow during this period, corresponding to a large decrease in H escape fluxes during the declining phase of the dust storm. Here we present airglow measurements of CO Cameron band and CO<sub>2</sub> UV Doublet emission in the thermosphere from 2007 data gathered by the SPICAM (SPECTROSCOPY FOR THE INVESTIGATION OF THE CHARACTERISTICS OF THE ATMOSPHERE OF MARS) instrument on Mars Express. During the dust storm period in late 2007, lower atmospheric temperatures were enhanced, pushing the thermosphere to higher altitudes. At the same time, more intense UV heating resulting from the perihelion passage of Mars resulted in extended upper atmospheric scale heights. We compare our retrievals and techniques to previous studies, which found no enhancement in thermospheric scale heights during this period. In addition, we present extended analysis of SPICAM H corona brightnesses, covering years 2005–2010. Sparse coverage of southern summer in years other than 2007 limits the usefulness of the dataset for constraining the maximum escape rates. However, the dataset is still useful for estimating H loss in the relatively quiet northern summer.

### SPICAM Observations of the Thermosphere:

The SPICAM instrument on Mars Express is an imaging ultraviolet spectrograph that observes in the far and mid UV from 118–320nm (Bertaux et al. 2006). The instrument has a focus on stellar occultations, but also performs thermospheric limb scans and nadir observations. In limb mode, the instrument observes the characteristic airglow spectrum of Mars (Figure 1), whose most conspicuous features in the mid-UV are the CO Cameron bands (180–260 nm), and the CO<sub>2</sub> UV doublet (289 nm). These features reveal thermospheric properties, with the UV Doublet diagnosing the neutral CO<sub>2</sub> scale height and the CO Cameron bands reflecting a larger scale height that is a blend of thermal and non-thermal emission sources. Previous studies, including most recently Stiepen et al. (2015), have examined the time evolution of the scale height and peak altitude of these features. Unfortunately, the need to share observing time

with other Mars Express instruments, and difficulties in the instrument data analysis introduced by stray light and detector high voltage spikes often limit the useful dataset, making observations sparse. As a result, these previous studies have extracted limited information about the thermosphere, particularly in the 2007 dust storm year. In addition, these previous studies found no large differences between 2007 and other years, despite the presumably large influence of the dust storm. This motivates our independent study of the dataset.




**Figure 1:** Mid-UV airglow of Mars, adapted from Leblanc et al. (2006).

### Retrieving Thermospheric Properties

To extract thermospheric peak altitudes and scale heights from the dataset, we employ a multiple linear regression technique. In this method, individual reduced spectra are fit by scaling input model spectra by linear coefficients, and summing the models together. For model spectra we use a Cameron band model convolved and binned to the instrument resolution (Stevens et al. 2015), a solar spectrum extracted from nadir observations made by SPICAM, and Gaussians at the location of the UV Doublet and OI 297.2 nm feature. We also include a linear component to account for dark subtraction offsets and large-scale stray light features.

An example fitted spectrum is shown in Figure 2a. Our method does a reasonable job of reproducing the data, with offsets in the vicinity of 240–260nm potentially caused by poor knowledge of the solar stray light at this location. Applying this technique to multiple spectra from a given limb scan produces intensity altitude


profiles (Figure 2b), which can then be fit to extract a peak altitude and scale height.



**Figure 2:** (a) Example SPICAM spectrum and multiple linear regression fit, showing composite fit and fitted components. (b) Retrieved altitude profile of CO Cameron band emission for a single limb scan. Each point represents the Cameron band fit component to the average SPICAM spectrum for surrounding altitudes.

**Response of the Thermosphere in 2007:**

By applying our fit procedure across 2007, we produce peak altitudes and scale heights for both the Cameron bands and the CO<sub>2</sub><sup>+</sup> UVD, shown in Figure 3. We see an increase of the peak altitude by over 10 km, centered on southern summer solstice and declining approximately linearly on either side of the peak. Scale heights are more peaked, with a linear increase from 15km near L<sub>s</sub> 200 to 30km near L<sub>s</sub> 300, and a steep decline following the peak. These observations have implications for the structure and variability of the thermosphere and the upper atmospheric environment underlying the large variability in the H corona, which we will discuss more fully in our presentation.




**Figure 3:** Retrieved peak altitudes and scale heights of thermospheric emissions at Mars in 2007. Red band indicates expected range of scale heights, orange band covers time range of 2007 global dust storm.

**SPICAM Observations of H Corona Variation:**

Following methods presented in Chaffin et al 2014 [5], we extended analysis of the SPICAM coronal Lyman alpha brightness dataset across 2005-2010, with results shown in Figure 4.

Unfortunately, the SPICAM UV data do not provide good coronal coverage of southern summers in years besides 2007, with most observations in other years falling into northern summer. This likely provides a good estimate of the minimum escape fluxes occurring in a given Martian year, without constraining the maximum H loss experienced in southern summer.



**Figure 4:** SPICAM retrieved H escape fluxes for 2005-2010. Panels show Martian dust opacity, H escape fluxes retrieved from SPICAM Lyman alpha profiles, Mars-Sun distance, and solar decametric radio flux at 1AU, an indicator of solar activity. Escape rates are highest in 2007 during Mars southern summer / perihelion, with lower rates at other times. Lack of coverage during southern summers other than 2007 limits the usefulness of the dataset in constraining the peak escape rates, but may provide a good estimate of escape during northern summer quiescent periods.

**References:**

[1] Bertaux, J.-L., et al.: SPICAM on Mars Express: Observing modes and overview of UV spectrometer data and scientific results, JGR, Vol. 111, E10S90, 2006.  
 [2] Stiepen, A., et al.: Mars thermospheric scale height: CO Cameron and CO<sub>2</sub><sup>+</sup> dayglow observations from Mars Express, Icarus, Vol. 245, pp. 295-305, 2015.  
 [3] Leblanc, F., et al.: Martian dayglow as seen by the SPICAM UV spectrograph on Mars Express, JGR, Vol. 111, E09S11, 2006.  
 [4] Stevens, M. H., et al.: New observations of molecular nitrogen in the Martian upper atmosphere by IUVS on MAVEN, GRL, Vol. 42, pp. 9050-9056, 2015.

[5] Chaffin, M.S., et al.: Unexpected variability of Martian hydrogen escape, *GRL*, Vol. 41, pp 314-320, 2014.