

Discovery of a New Tidal Disruption Event Candidate from the 2XMM Catalog

Dacheng Lin(dlin@cesr.fr)¹, Eleazar R. Carrasco², Dirk Grupe³, Natalie A. Webb¹, Didier Barret¹, & Sean A. Farrell⁴



1. Institut de Recherche en Astrophysique et Planétologie, France; 2. Gemini Observatory, Southern Operation Center, Chile;
3. Pennsylvania State University, USA; 4. The University of Sydney, Australia



1. Introduction

Stars approaching supermassive black hole (SMBHs) can be tidally disrupted and subsequently accreted. Such events provide a unique way to find and study dormant SMBHs believed to reside in many galaxies. About a dozen of such event candidates are found, with peak spectra typically soft ($\sim 0.04\text{--}0.1$ keV) and luminosities up to 10^{44} erg/s. We report our discovery of a new candidate, with two ultrasoft X-ray spectra of unprecedented quality near the peak (Lin et al. 2011, ApJ in press, arXiv:1106.0744).



Fig 1. Artist's sketch of the tidal disruption of a star around a SMBH

(3) **Ultrasoft X-ray spectra.** Both XMM-Newton spectra can be described by a strong thermal disk plus a weak power law (Figure 5). The disk temperatures ($\sim 0.07\text{--}0.1$ keV) and luminosities are consistent with $L \propto T^4$, similar to BH X-ray binaries in the thermal state but for a SMBH of $\sim 10^6 M_\odot$.




Fig 6. The disk flux versus the temperature. The solid ($L \propto T^4$), dotted ($L \propto T^3$), and dashed ($L \propto T^5$) lines are for guidance.




Fig 5. The unfolded pn spectrum and the fit residuals (black/red/green for pn/MOS1/MOS2, respectively), using a multicolor disk model (dotted line) plus a power law (dashed line).

(4) **Fast X-ray variability.** Both XMM-Newton observations show large variability, with rms values about 21% on timescales of hours (Figure 7). The disk temperature is found to increase with the flux, implying a possible explanation for the fast variability: it is due to the fast variation of the accretion rate caused by shocks during tidal disruption of the star.




Fig 7. Left panels: the pn 0.2-2.0 keV light curves. Right panels: the power density spectra. The dotted line is the fit with a power law plus a constant Poisson level (the solid line).

2. A New Candidate: 2XMMi J184725.1-631724

(1) Lying toward the center of a galaxy ($z=0.0353$, Figure 2), which is inactive: no clear optical emission lines but typical stellar absorption features (Figure 3)




Fig 2. The V-band image of the galaxy. The green plus is the galaxy center from the profile fit. The red circle is the $3\text{-}\sigma$ error of the X-ray position.

Fig 3. The optical spectrum of the galaxy from the Gemini South Telescope four years after the flare.

(2) **An X-ray transient source.** It was detected serendipitously in two XMM-Newton observations (XMM1 and XMM2 in Figure 4), but not in a ROSAT observation in 1992 or a Swift observation in 2011.




Fig 4. The long-term luminosity curve. Arrows are $3\text{-}\sigma$ upper limits. The dashed and solid lines are two theoretic decay curves.

3. Conclusion

2XMMi J184725.1-631724 is an ultrasoft X-ray transient, with peak luminosity near 10^{44} erg/s and lying toward the center of an inactive galaxy, thus probably a tidal disruption event. From its two ultrasoft X-ray observations near the peak of unprecedented quality, we gain more insights into such events:

- Near the flare peak, the disk luminosity appears to follow $L \propto T^4$, similar to BH X-ray binaries in the thermal state but for a SMBH;
- Large variability on timescales of hours are seen, probably due to fast variations in the mass accretion rate caused by shocks during tidal disruption of the star.