

Max-Planck-Institut für extraterrestrische Physik /







Screening and validation of EXTraS data products


www.extras-fp7.eu

S. Carpano¹, F. Haberl¹, A. De Luca², A. Tiengo³, G. Israel⁴, G. Rodriguez⁴, A. Belfiore², S. Rosen⁵, A. Read⁵, J. Wilms⁶, A. Kreikenbohm⁶, D. Law-Green⁵

¹MPE, Garching ²INAF/IASF Milano ³IUSS Pavia ⁴INAF/OA Roma ⁵University of Leicester ⁶ECAP Univ. Erlangen-Nürnberg




The EXTraS project (Exploring the X-ray Transient and variable Sky) is aimed at fully exploring the serendipitous content of the XMM-Newton EPIC database in the time domain. The project is funded within the EU/FP7-Cooperation Space framework and is carried out by a collaboration including INAF (Italy), IUSS (Italy), CNR/ IMATI (Italy), University of Leicester (UK), MPE (Germany) and ECAP (Germany). The several tasks consist in characterise aperiodic variability for all 3XMM sources, search for short-term periodic variability on hundreds of thousands sources, detect new transient sources that are missed by standard source detection and hence not belonging to the 3XMM catalogue, search for long term variability by measuring fluxes or upper limits for both pointed and slew observations, and finally perform multiwavelength characterisation and classification. Screening and validation of the different products is essential in order to reject flawed results, generated by the automatic pipelines. We present here the screening tool we developed in the form of a Graphical User Interface and our plans for a systematic screening of the different catalogues.



Visualisation of PPS

products The GUI displays images like background, exposure maps but also scientific images in one or several bands (including RGB images), plots of the background light curve, changes in the attitude, light curves and spectra of brightest sources, and prints information about the observation.



Visualisation of the WP2 products

(aperiodic variability) The GUI displays light curves (with different binning times), Bayesian Block light curves, Cumulative Rate Functions, Fast Fourier Transforms and Energy spectra. It also allows the user to produce its own light curves using different settings.

Visualisation of WP4 products (transient sources)

The GUI displays images of the transient candidates in the relevant time interval, with exposure and background maps, prints the detection parameters and allows the user to produce its own light curve using different settings.


SHOW ALL TRANS	ID_I	NST-0: all		M1, 3: M2	0.5], 2: [0.5	-1], 3:	[1-2], 4: [2:	4.5], 5: [4.5-12]								
Select Trans: Trans 1 (TF=1)		ID_INST	ID_BAND	SCTS (sta)	DET_ML	D/T	DOP_MAP		~	0	0	+	a		v	
	1	0	0	72.4684	73.7204	0.0	nan			· ·	× .	•				
SHOW MAPS	2	1	0	37.9817	44.6805	0.0	nan							transl		


Description of the GUI

The GUI allows a quick visualisation of the XMM-Newton PPS files, EXTraS data products and optical dss image of the FOV. Written in Python, it has a plot/graphic window and an interface to ds9 for image display.

LTV SRCID: or enter LTV RA, DEC, RAD: RA (d):	OBSERV ID: 010406010 RA (d): 82.8333349 DEC (d): -65.960555	S OPN O	M1 M2	0 + 🧭	8 🖩 🗸	
DEC (d):	Keep RA,DEC			53112.9-860706 - peak 1 c	LMC deep field cd00 Bin time:	1.367 s
RAD ("): 10	SHOW TRANSIENT DA	TA				
	SHOW PERIODIC SRC	s		Best Period Resolution	1: 13.6454997944781 s n: 0.131E-03 s	
\$	PLOT USER WP2 LC	Select EXP ID):		6	
Done	SHOW DSS IMAGE	PNS001 M2S003	_ M15002			
Amplitude= 3.95951e-01 Proba= 1.18769e-22 Power= 1.36154e+02 Ffr= 2815 0 0 0	PLOT ATT		DBS INFO			
Colored Bardin Reserved	Scargle period search:	Scargle parameters:				
Select Periodic Source:	Use source files:	PIMIN (eV): 20	00		11	
select Periodic Source:	o use source mes:				A	mound
	M1 Mleps/ev v	PIMAX (eV)	0000	المحطان فالرهول فالجادوا حسرتين الداء	when a server	
[src001 :]	M1 Mlepx/ev y M2 M2epx/ev y	PE_MIN (s):		-0.06 -0.04 -0.02	0 0.02 0.04	
SHOW IMAGES	✓ M1 Mlepx/ev v ✓ M2 M2epx/ev v ✓ PN PNeps/evt v	PE_MIN (s): 10 PE_MAX (s): 50		-0.06 -0.04 -0.02		
Select run: epx/evt1 :	M1 Mlepx/ev y M2 M2epx/ev y	PE_MIN (s): IN PE_MAX (s): 24 TL_MIN (ks), orig. LC:		-0.06 -0.04 -0.02	0 0.02 0.04	0.06
Select run: epx/evt1 : SHOW DPS	✓ M1 Mlepx/ev v ✓ M2 M2epx/ev v ✓ PN PNeps/evt v	PE_MIN (s): IX PE_MAX (s): 54 TI_MIN (ks), orlg. LC: TI_MAX (ks), orig. LC:		-0.06 -0.04 -0.02	0 0.02 0.04 Period (s) - Offset	0.06
SHEDD1 : SHOW IMAGES Select run: epx/evt1 : SHOW DPS SHOW PSD	✓ M1 M1epx/ev y ✓ M2 M2epx/ev y ✓ M1 Miepx/ev y ✓ M2 Miepx/ev y ✓ M1 Miepx/ev y ✓ M1 Miepx/ev y	PE_MIN (s): IN PE_MAX (s): 74 TI_MIN (ks), orig. LC: TI_MAX (ks), orig. LC: Nb periods: 74		-0.06 -0.04 -0.02	0 0.02 0.04 Period (s) - Offset	0.06
Select run: epx/evt1 : SHOW DPS	M1 Mlepx/ev y M2 M2apx/ev y M2 MPeps/ev y Use background light curves: PN5001 M15002	PE_MIN (s): IX PE_MAX (s): 54 TI_MIN (ks), orlg. LC: TI_MAX (ks), orig. LC:		-0.06 -0.04 -0.02	0 0.02 0.04 Period (s) - Offset	0.06

Visualisation of WP3 products (periodic variability) The GUI displays graphs of the Discrete Period Search, Power Spectra Density, folded light curve and Epoch Folding search around period. The user can also perform its own period search via Lomb Scargle Periodograms, using different settings.





(long term variability) The GUI displays long term light curves using the WP5 catalogue which contains fluxes and upper limits from pointed observations and slew surveys. It prints variability parameters and shows images of the sources in the different observations.

Visualisation of WP5 products

Plans for systematic screening of the different catalogues

For each WP it is planned to look for the most variable sources according to selected criteria and filtering. The GUI will then be used to check if the variability is real or spurious (because of bad data quality or imperfections in the automatic pipelines). After a careful analysis, feedback will be given to the different WPs to improve the algorithm and/or data products quality.